arXiv:1304.2298v1 [math.GT] 8 Apr 2013

A DISCRETENESS CRITERION FOR GROUPS CONTAINING SCREW TRANSLATIONS

VIVEKA ERLANDSSON AND SAEED ZAKERI

Abstract. This note will prove a discreteness criterion for groups of orientation-preserving isometries of the 4-dimensional hyperbolic space which contain an irrational screw translation. It can be viewed as a generalization of the well-known results of Shimizu-Leutbecher and Jørgensen in dimensions 2 and 3, and is in the same spirit as Waterman's inequality for higher dimensions. Unlike his algebraic method, the argument presented here is geometric and yields a sharper asymptotic bound.

1. The result

We will need a few basic facts about the hyperbolic space and its isometries. The material is standard and can be found in several references such as [2], [6], and [9], to name a few.

Throughout we will use the upper half-space model for the 4-dimensional real hyperbolic space:

$$\mathbb{H}^4 = \{x = (v,t) : v \in \mathbb{R}^3, t > 0\}.$$

The extended boundary $\partial\mathbb{H}^4 = \mathbb{R}^3 \cup \{\infty\}$ is homeomorphic to the 3-sphere, with the closure $\overline{\mathbb{H}}^4 = \mathbb{H}^4 \cup \partial\mathbb{H}^4$ homeomorphic to the closed 4-ball. The hyperbolic metric $(dv^2 + dt^2)/t^2$ on $\mathbb{H}^4$ induces the distance $\rho(\cdot,\cdot)$ which satisfies

$$\cosh(\rho(x,\hat{x})) = 1 + \frac{\|x-\hat{x}\|^2}{2t\hat{t}} \qquad (x,\hat{x} \in \mathbb{H}^4). \tag{1}$$

Here $\|\cdot\|$ is the Euclidean norm in $\mathbb{R}^4$ and $t, \hat{t}$ are the last coordinates of $x, \hat{x}$.

We denote by $\mathrm{Isom}^+(\mathbb{H}^4)$ the group of orientation-preserving isometries of $\mathbb{H}^4$ with respect to hyperbolic metric. Every element of $\mathrm{Isom}^+(\mathbb{H}^4)$ extends continuously to a Möbius map acting on $\partial\mathbb{H}^4$. Conversely, the "Poincaré extension" of every Möbius map of $\partial\mathbb{H}^4$ is an element of $\mathrm{Isom}^+(\mathbb{H}^4)$. It follows that $\mathrm{Isom}^+(\mathbb{H}^4)$ is canonically isomorphic to the group Möb(3) of orientation-preserving Möbius maps acting on the 3-sphere. For each $g \in \mathrm{Isom}^+(\mathbb{H}^4)$, the fixed point set $\mathrm{Fix}(g) = \{x \in \overline{\mathbb{H}}^4 : g(x) = x\}$ is non-empty. A non-identity g is ***elliptic*** if $\mathrm{Fix}(g)$ intersects $\mathbb{H}^4$, ***loxodromic*** if

---

Date: April 9, 2013.

2010 *Mathematics Subject Classification.* 22E40, 30F40.

$\mathrm{Fix}(g)$ consists of two distinct points on $\partial\mathbb{H}^4$, and ***parabolic*** if $\mathrm{Fix}(g)$ consists of a unique point on $\partial\mathbb{H}^4$. Every parabolic isometry g is conjugate to the normal form

$$(v,t) \mapsto (Av+a,t) \qquad (v \in \mathbb{R}^3, t>0) \tag{2}$$

fixing $\infty \in \partial\mathbb{H}^4$, where $A \in \mathrm{SO}(3)$ and $a \in \mathbb{R}^3$ is non-zero with $Aa=a$. The conjugacy class of A in $\mathrm{SO}(3)$ is uniquely determined by g. We call g a ***pure translation*** if $A=I$, and a ***screw translation*** if $A \neq I$. A screw translation is ***rational*** if A has finite order, and is ***irrational*** otherwise. These correspond to the cases where the angle of rotation of A about its axis is a rational or irrational multiple of 2π.

Consider a discrete group $\Gamma \subset \mathrm{Isom}^+(\mathbb{H}^4)$ which acts freely on $\mathbb{H}^4$ and contains a parabolic element which fixes, say, $\infty \in \partial\mathbb{H}^4$. The stabilizer

$$\Gamma_\infty = \{g \in \Gamma : g(\infty)=\infty\}$$

is then a maximal parabolic subgroup of Γ. For $\varepsilon>0$, define the ***Margulis region*** associated with Γ_∞ by

$$T(\Gamma_\infty) = \big\{x \in \mathbb{H}^4 : \rho(g(x),x) < \varepsilon \text{ for some } g \in \Gamma_\infty \smallsetminus \{\mathrm{id}\}\big\}. \tag{3}$$

It is not hard to see that $T(\Gamma_\infty)$ is a non-empty domain in $\mathbb{H}^4$ having ∞ on its boundary. Furthermore, when ε is small (less than the Margulis constant of $\mathbb{H}^4$), $T=T(\Gamma_\infty)$ is ***precisely invariant*** under Γ_∞ in the sense that

$$\begin{cases} g(T)=T & \text{if } \ g \in \Gamma_\infty \\ g(T) \cap T = \emptyset & \text{if } \ g \in \Gamma \smallsetminus \Gamma_\infty \end{cases}$$

(see [3]). When Γ_∞ consists only of pure translations or rational screw translations, $T(\Gamma_\infty)$ contains a precisely invariant horoball based at ∞. But when there is an irrational screw translation in Γ_∞, the Margulis region $T(\Gamma_\infty)$ cannot contain a horoball. In fact, examples constructed by Apanasov [1] and Ohtake [5] show that such groups may fail to have any precisely invariant horoball. In this case, the geometry of the Margulis region becomes relevant in understanding the corresponding parabolic end of the quotient manifold $\mathbb{H}^4/\Gamma$.

To describe this geometry more explicitly, following [8] we use the coordinates (r,θ,z,t) in $\mathbb{H}^4$, where (r,θ,z) are the cylindrical coordinates of $\mathbb{R}^3$ and $t>0$. We assume $\theta \in \mathbb{R}/\mathbb{Z}$, which amounts to measuring the polar angle in full turns rather than multiples of 2π. It follows from the normal form (2) that every parabolic isometry is conjugate to the map

$$g_\alpha : (r,\theta,z,t) \mapsto (r,\theta+\alpha,z+1,t) \tag{4}$$

for a unique angle $\alpha \in \mathbb{R}/\mathbb{Z}$. When the stabilizer Γ_∞ contains an irrational screw translation, it is necessarily a cyclic group [3], so we may assume without loss of generality that Γ_∞ is generated by the map g_α for some irrational α. As the Margulis region $T(\Gamma_\infty)$ is now independent of the rest of the group Γ, and to emphasize its sole dependence on α, we denote it by T_α. It follows from the definition (3) that

$$T_\alpha = \bigcup_{j=1}^{\infty} T_{\alpha,j}, \tag{5}$$

where

$$T_{\alpha,j} = \{x \in \mathbb{H}^4 : \rho(g_\alpha^{\circ j}(x), x) < \varepsilon\}.$$

Note that we have taken the union over positive j only since $T_{\alpha,j} = T_{\alpha,-j}$. By (1), the condition $(r, \theta, z, t) \in T_{\alpha,j}$ is equivalent to

$$\frac{\|(r, \theta + j\alpha, z + j, t) - (r, \theta, z, t)\|^2}{2t^2} < \cosh(\varepsilon) - 1$$

or

$$\frac{4\sin^2(\pi j\alpha)\ r^2 + j^2}{2t^2} < \cosh(\varepsilon) - 1.$$

Define the sequence of functions $u_{\alpha,j} : [0, \infty) \to \mathbb{R}$ by

$$u_{\alpha,j}(r) = c(\varepsilon)\sqrt{4\sin^2(\pi j\alpha)\ r^2 + j^2}, \tag{6}$$

where $c(\varepsilon) = 1/\sqrt{2\cosh(\varepsilon) - 2}$. Then,

$$T_{\alpha,j} = \{(r, \theta, z, t) \in \mathbb{H}^4 : t > u_{\alpha,j}(r)\}.$$

If we define the ***boundary function*** $\mathscr{B}_\alpha : [0, \infty) \to \mathbb{R}$ by

$$\mathscr{B}_\alpha(r) = \inf_{j \geq 1} u_{\alpha,j}(r),$$

it follows from (5) that

$$T_\alpha = \{(r, \theta, z, t) \in \mathbb{H}^4 : t > \mathscr{B}_\alpha(r)\}.$$

It is easy to see that $\mathscr{B}_\alpha$ is strictly increasing and piecewise smooth. Moreover, the asymptotic behavior of $\mathscr{B}_\alpha(r)$ as $r \to \infty$ turns out to depend on the arithmetical properties of α determined by its continued fraction expansion (see [8] and [3]). In a recent work, we have investigated this dependence to prove the following result:

Theorem 1 ([3])**.** *For every irrational $\alpha \in \mathbb{R}/\mathbb{Z}$, the boundary function $\mathscr{B}_\alpha$ of the Margulis region T_α satisfies the asymptotically universal upper bound*

$$\mathscr{B}_\alpha(r) \leq \text{const.}\ r^{1/2} \qquad \textit{for large } r.$$

If α is Diophantine of exponent $\nu \geq 2$, then $\mathscr{B}_\alpha$ satisfies the lower bound

$$\mathscr{B}_\alpha(r) \geq \text{const.}\ r^{1/(2\nu-2)} \qquad \textit{for large } r.$$

The upper bound is asymptotically universal in the sense that the constant involved is independent of α. In fact, a quantitative version of this result shows that

$$\sup_{\alpha} \limsup_{r\to\infty} \frac{\mathscr{B}_\alpha(r)}{\sqrt{r}} < 1{,}000.$$

On the other hand, there are irrationals α of Liouville type for which $\mathscr{B}_\alpha$ has arbitrarily slow growth over long intervals [3].

In this note, we will use the geometry of the Margulis region to prove a discreteness criterion for subgroups of $\mathrm{Isom}^+(\mathbb{H}^4)$ which contain irrational screw translations. Our result can be viewed as a generalization of the well-known results of Shimizu-Leutbecher [7] and Jørgensen [4] in dimensions 2 and 3 which give a uniform bound on the radii of the isometric spheres of elements in a discrete group which move the parabolic fixed point ∞. In higher dimensions, Ohtake [5] has shown that no such uniform bound exists in the presence of an irrational screw translation. In [10], Waterman obtains a bound on the radii of the isometric spheres as a function of the Euclidean distance that their centers move under the screw translation. Our result similarly bounds the radii in terms of the centers of the isometric spheres.

For each $h \in \mathrm{Isom}^+(\mathbb{H}^4)$ with $h(\infty) \neq \infty$, denote the isometric circle of h by I_h, the Euclidean radius of I_h by R_h, and the r-coordinate of the center of I_h by r_h.

Theorem 2 (Discreteness Criterion)**.** *Let* $h \in \mathrm{Isom}^+(\mathbb{H}^4)$ *with* $h(\infty) \neq \infty$. *Suppose the subgroup* $\langle h, g_\alpha\rangle \subset \mathrm{Isom}^+(\mathbb{H}^4)$ *is discrete for some irrational* $\alpha \in \mathbb{R}/\mathbb{Z}$. *Then*

$$R_h \leq \big(\mathscr{B}_\alpha(r_h)\ \mathscr{B}_\alpha(r_{h^{-1}})\big)^{1/2}.$$

In particular, we have the asymptotically universal bound

$$R_h \leq \text{const.}\ (r_h\, r_{h^{-1}})^{1/4} \qquad \textit{for large } r_h, r_{h^{-1}}.$$

This asymptotic bound turns out to be substantially better than the one in [10]; see the discussion at the end of the next section.

2. The Proof

Let us recall a few basic facts about isometric circles. Consider a map $h \in \mathrm{Möb}(3)$ with $h(\infty) \neq \infty$, and let $v = h^{-1}(\infty)$, $v' = h(\infty)$. It is not hard to check that h maps each Euclidean sphere $S(v, R) = \{w \in \mathbb{R}^3 : \|w - v\| = R\}$ to a Euclidean sphere $S(v', R')$. Here $R \mapsto R'$ is a decreasing function, with $\lim_{R\to 0} R' = \infty$ and $\lim_{R\to\infty} R' = 0$. Hence there is a unique radius $R_h > 0$ such that $h(S(v, R_h)) = S(v', R_h)$. We call $S(v, R_h)$ the ***isometric sphere*** of h and denote it by I_h. The map $h : I_h \to I_{h^{-1}}$ is a Euclidean isometry, and $R_h = R_{h^{-1}}$. Moreover, h maps the interior (resp. exterior) of I_h to the exterior (resp. interior) of $I_{h^{-1}}$.

If $h \in \mathrm{Isom}^+(\mathbb{H}^4)$ with $h(\infty) \neq \infty$, we define the isometric sphere I_h as the convex hull of the isometric sphere of the boundary map $h|_{\partial\mathbb{H}^4} \in \mathrm{Möb}(3)$, that is, the upper



Figure 1. *Proof of Theorem 2*

hemisphere in $\mathbb{H}^4$ with the same center and radius as those of $I_{h|_{\partial\mathbb{H}^4}}$. The center of I_h lies on $\partial\mathbb{H}^4$, so it has coordinates $(v_h, 0) \in \mathbb{R}^3 \times \{0\}$. We denote by r_h the distance from v_h to the z-axis in $\mathbb{R}^3$. In terms of our coordinates (r, θ, z, t) of $\mathbb{H}^4$, this is just the r-coordinate of the center of I_h. The north pole N_h of the isometric circle I_h has coordinates (v_h, R_h). Since $h(v_h, 0) = \infty$ and $h(\infty) = (v_{h^{-1}}, 0)$, h carries the vertical geodesic from $(v_h, 0)$ to ∞ isometrically to the vertical geodesic from $(v_{h^{-1}}, 0)$ to ∞. It follows in particular that $h(N_h) = N_{h^{-1}}$.

Proof of Theorem 2. For convenience we drop the subscript h from our notations and denote the quantities associated with h^{-1} with a minus sign. There is nothing to prove if $R \leq \mathscr{B}_\alpha(r)$ and $R \leq \mathscr{B}_\alpha(r^-)$, so let us assume that $R > \mathscr{B}_\alpha(r)$ (the case $R > \mathscr{B}_\alpha(r^-)$ is similar). The vertical geodesic segment σ from the north pole $N = (v, R)$ down to $(v, \mathscr{B}_\alpha(r))$ lies in the Margulis region T_α and has hyperbolic length $\log(R/\mathscr{B}_\alpha(r))$ (see Fig. 1). The image $h(\sigma)$ is the vertical geodesic segment from $N^- = (v^-, R)$ up to (v^-, t) for some t, and has the same length as σ. It follows that $\log(t/R) = \log(R/\mathscr{B}_\alpha(r))$, or $R^2 = \mathscr{B}_\alpha(r)t$. Let $\Gamma = \langle h, g_\alpha \rangle$ and let $\Gamma_\infty = \langle g_\alpha \rangle$ be the stabilizer of ∞. Since T_α is precisely invariant under the action of Γ_∞ and $h \notin \Gamma_\infty$, the image $h(\sigma)$ must be disjoint from T_α. This shows $t < \mathscr{B}_\alpha(r^-)$, and proves the required inequality $R^2 < \mathscr{B}_\alpha(r)\mathscr{B}_\alpha(r^-)$.

The asymptotically universal bound $R \leq \text{const.}\,(r\,r^-)^{1/4}$ follows from this and Theorem 1. □

A comment on Theorem 2 is in order. Waterman's result in [10] is the inequality

$$R_h \leq K_\alpha \,\|g_\alpha(v_h) - v_h\|^{1/2}\, \|g_\alpha(v_{h^{-1}}) - v_{h^{-1}}\|^{1/2},$$

where K_α is a constant depending on α. He derives this via a purely algebraic argument based on Clifford matrix representation of Möbius maps, and assuming $\operatorname{dist}(\alpha, \mathbb{Z})$ is small. Replacing g_α by the iterate $g_\alpha^{\circ j}$, and using the distance formula

(1) and the definition of the functions $u_{\alpha,j}$ in (6), one can re-write the above inequality as

$$R_h \leq \frac{K_\alpha}{c(\varepsilon)} \left(u_{\alpha,j}(r_h)\ u_{\alpha,j}(r_{h^{-1}})\right)^{1/2} \qquad (j \geq 1). \tag{7}$$

Since each $u_{\alpha,j}$ is asymptotically linear in r, for large $r_h, r_{h^{-1}}$ these bounds take the form

$$R_h \leq \text{const.}\ (r_h r_{h^{-1}})^{1/2}.$$

Our geometric approach, in effect, takes the infimum over all j of the upper bounds in (7), which yields the improved 4th-root asymptotic bound. It is not hard to construct 2-generator groups of isometries of $\mathbb{H}^4$ containing g_α that are non-discrete by Theorem 2, for which Waterman's criterion remains inconclusive.

References

[1] B. Apanasov, *Cusp ends of hyperbolic manifolds*, Ann. Global Analysis and Geometry, **3** (1985) 1-11.

[2] R. Benedetti and C. Petronio, *Lectures on Hyperbolic Geometry*, Springer, 2003.

[3] V. Erlandsson and S. Zakeri, *On Margulis cusps of hyperbolic 4-manifolds*, submitted, 2013.

[4] T. Jørgensen, *On discrete groups of Möbius transformations*, Amer. J. Math. **98** (1976) 739-749.

[5] H. Ohtake, *On discontinuous subgroups with parabolic transformations of the Möbius groups*, J. Math. Kyoto Univ., **25** (1985) 807-816

[6] J. Ratcliffe, *Foundations of hyperbolic manifolds*, Springer, 1994.

[7] H. Shimizu, *On discontinuous groups operating on the product of the upper half planes*, Ann. of Math. **77** (1963) 33-71.

[8] P. Susskind, *The Margulis region and continued fractions*, Complex manifolds and hyperbolic geometry (Guanajuato, 2001), Contemp. Math., **311**, Amer. Math. Soc., Providence, RI, (2002) 335-343.

[9] W. Thurston, *Three-dimensional geometry and topology, Vol. 1*, Princeton University Press, 1997.

[10] P. Waterman, *Möbius transformations in several dimensions*, Adv. in Math., **101** (1993) 87-113.

V. Erlandsson, Department of Mathematics, Graduate Center of CUNY, New York

E-mail address: `viveca@gmail.com`

S. Zakeri, Department of Mathematics, Queens College and Graduate Center of CUNY, New York

E-mail address: `saeed.zakeri@qc.cuny.edu`